

# RCFT extensions of $W_{1+\infty}$ in terms of bilocal fields

L.S. Georgiev <sup>\*</sup> I.T. Todorov <sup>†</sup>

*Institute for Nuclear Research and Nuclear Energy  
Tsarigradsko Chaussee 72, BG-1784 Sofia, Bulgaria*

## Abstract

The rational conformal field theory (RCFT) extensions of  $W_{1+\infty}$  at  $c = 1$  are in one-to-one correspondence with 1-dimensional integral lattices  $L(m)$ . Each extension is associated with a pair of oppositely charged “vertex operators” of charge square  $m \in \mathbb{N}$ . Their product defines a bilocal field  $V_m(z_1, z_2)$  whose expansion in powers of  $z_{12} = z_1 - z_2$  gives rise to a series of (neutral) local quasiprimary fields  $V^l(z, m)$  (of dimension  $l + 1$ ). The associated bilocal exponential of a normalized current generates the  $W_{1+\infty}$  algebra spanned by the  $V^l(z, 1)$  (and the unit operator). The extension of this construction to higher (integer) values of the central charge  $c$  is also considered.

Applications to a quantum Hall system require computing characters (i.e., chiral partition functions) depending not just on the modular parameter  $\tau$ , but also on a chemical potential  $\zeta$ . We compute such a  $\zeta$  dependence of orbifold characters, thus extending the range of applications of a recent study of affine orbifolds.

---

<sup>\*</sup>E-mail: lgeorg@inrne.acad.bg

<sup>†</sup>E-mail: toodorov@inrne.acad.bg

# 1 Introduction

The unitary positive energy representations of  $W_{1+\infty}$  [KR1] correspond to positive integer central charges  $c \in \mathbb{N}$ . A standard tool in their study [KR1, KR2] [FKRW] [AFMO] is the embedding of all irreducible modules for a given  $c$  into the Fock space of  $c$  pairs of free Weyl fermion fields  $\psi_i^*(z)$ . It has been used in [BGT] to develop a conformal (quantum) field theory (CFT) approach to  $W_{1+\infty}$  based on the expansion of the bilocal field

$$V_1(z_1, z_2) = \sum_{i=1}^c \left( \psi_i^*(z_1) \psi_i(z_2) - \frac{1}{z_{12}} \right), \quad z_{12} = z_1 - z_2 \quad (1.1)$$

into local quasiprimary fields.  $V_1$  is invariant under  $\mathrm{SO}(2c)(\supset \mathrm{U}(1)^{\times c})$  gauge transformations in the  $2c$ -dimensional space of  $\psi_i$  and  $\psi_i^*$  (it is, in particular,  $\mathrm{U}(1)^{\times c}$  neutral). The index 1 on  $V$  indicates the charge square of  $\psi_i$  (and  $\psi_i^*$ ) which coincides with the (total) conformal dimension of  $V_1$ : there is a representation  $U(\rho)$  of the multiplicative group of positive reals ( $\rho > 0$ ) in the state space of our CFT such that

$$U(\rho)V_1(z_1, z_2)U(\rho)^{-1} = \rho V_1(\rho z_1, \rho z_2) \quad (\text{and } U(\rho)|0\rangle = |0\rangle) \quad (1.2)$$

( $|0\rangle$  representing the unique zero-energy state, the *vacuum* of the theory). The chiral algebra  $W_c$  of  $c$  Weyl fermions (and their conjugates) corresponds to the simplest orthogonal lattice,  $\mathbb{Z}^c$ , of admissible charges:  $W_c = \mathcal{A}(\mathbb{Z}^c)$ . In general, the abelian compact Lie group  $\mathrm{U}(1)^{\times c}$  defines a lattice in  $\mathfrak{u}(1)^{\otimes c}$  of elements  $\omega$  satisfying  $e^{2\pi i \omega} = 1$  (where 1 stands for the group unit). Equipped with a positive definite integral quadratic form,  $(\omega|\omega') \in \mathbb{Z} \quad |\omega|^2 = (\omega|\omega) \in \mathbb{N}$  for  $\omega \neq 0$ ,  $L$  gives rise to a chiral algebra  $\mathcal{A}(L)$  whose positive energy (localizable) representations give rise to an RCFT (see, e.g., Sec. 1 of [KT]). Singling out a current  $H(z)$  in the Kac-Moody algebra  $\hat{\mathfrak{u}}(1)^{\otimes c}$  we can view  $\mathcal{A}(L)$  as an RCFT extension of the level  $c$   $W_{1+\infty}$  algebra which contains  $H$ . We define a bilocal field  $V_m(z_1, z_2)$ , expressed as a bilocal exponential in  $H$  - see (2.17) (generalizing (1.1) to any positive integer charge square  $m$ ) and establish for it an *operator product expansion* (OPE) of quasiprimary fields  $V^l(z, m)$  of dimension  $l+1$  ( $l = 0, 1, \dots$ ). We prove, however, that only for  $m = 1$ , in which case the fields  $V^i(z) = V^l(z, 1)$  are expressed in terms of an orthonormal set of  $\mathfrak{u}(1)$  currents  $J^i$ ,

$$\langle J^i(z_1) J^j(z_2) \rangle = \delta_{ij} z_{12}^{-2} \quad i, j = 1, \dots, c, \quad (1.3)$$

do the  $V^l$  (and the unit operator) close an infinite dimensional Lie algebra (which coincides with the level  $c$   $W_{1+\infty}$ ). We work out the details, establishing exact OPE formulae, in Sec. 2 for the case  $c = 1$  and indicate the extension to arbitrary  $c \in \mathbb{N}$  in Sec. 3.

## 2 Rational extensions of $W_{1+\infty}$ for $c = 1$ . OPE of quasiprimary bilocal fields

### 2.1 Local field basis for $W_{1+\infty}$ in terms of Schur polynomials of a $u(1)$ current and its derivatives

The product of two oppositely charged fields  $\phi^\pm(z)$  of charge square  $m$  can be expressed as a normal ordered exponential of a  $U(1)$  current [DFZ]:

$$\phi^+(z_1)\phi^-(z_2) = z_{12}^{-m} : \exp \left\{ \int_{z_2}^{z_1} H(z) dz \right\} :, \quad z_{12} = z_1 - z_2. \quad (2.1)$$

(The subtractions involved in the definition of normal ordering can be formally described by dividing the exponential by its vacuum expectation value.) The normalization of the “Cartan current”  $H$  can be read off the singular terms of the OPE:

$$H(z_1)H(z_2) \sim \frac{m}{z_{12}^2}, \quad H(z_1)\phi^\pm(z_2) \sim \frac{\pm m}{z_{12}} \phi^\pm(z_2). \quad (2.2)$$

For a  $c = 1$  theory the stress energy tensor  $T(z)$  is expressed in terms of  $H$  by the Sugawara formula:

$$T(z) = \frac{1}{2m} : H^2(z) : \quad (: H(z_1)H(z_2) : \equiv H(z_1)H(z_2) - \frac{m}{z_{12}^2}). \quad (2.3)$$

Its Laurent modes  $L_n$  generate the Virasoro algebra:

$$T(z) = \sum_{n \in \mathbb{Z}} L_n z^{-n-2} \quad ([L_n, L_{n'}] = (n - n')L_{n+n'} + \frac{c}{12}(n^3 - n)\delta_{n,-n'}, \quad c = 1). \quad (2.4)$$

A central result in [BGT] says that the bilocal field (2.1) is, for  $m = 1$ , a generating functional for a local field basis of the  $W_{1+\infty}$  algebra. In fact, it then coincides (up to a  $c$ -number term) with (1.1) for central charge  $c = 1$ . We shall reformulate this result using the  $U(1)$  current alone (without reference to the free Weyl field  $\psi$ ) thus extending it to the more general current algebra framework.

Denote the normalized  $U(1)$  current  $\frac{1}{\sqrt{m}}H(z)$  by  $J^0(z)$ . Using the Taylor expansion of the integral in the exponential in (2.1),

$$\int_{z_2}^{z_1} J^0(z) dz = \sum_{l=0}^{\infty} \frac{z_{12}^{l+1}}{(l+1)!} \partial_2^l J^0(z_2) \quad (\partial_2 = \frac{\partial}{\partial z_2}; \quad \langle J^0(z_1)J^0(z_2) \rangle_0 = z_{12}^{-2}) \quad (2.5)$$

we introduce the *Schur polynomials* (in the terminology of [KR3] and [BHY]) of  $J^0$  and its derivatives,

$$J^l(z) = l! :S_{l+1}(J^0(z), \frac{\partial J^0(z)}{2!}, \dots, \frac{\partial^l J^0(z)}{(l+1)!}):, \quad (2.6)$$

setting

$$V_1(z_1, z_2) = \frac{1}{z_{12}} \left( : \exp \left\{ \int_{z_2}^{z_1} J^0(z) dz \right\} : - 1 \right) = \sum_{l=0}^{\infty} \frac{z_{12}^l}{l!} J^l(z_2). \quad (2.7)$$

In general, a Schur polynomial of  $N$  variables is defined by

$$S_N(t_1, t_2, \dots, t_k, \dots) = \sum_{\substack{n_1, n_2, \dots, n_k, \dots \\ \sum_{k=1}^{\infty} k n_k = N}} \frac{t_1^{n_1} t_2^{n_2} \dots t_k^{n_k} \dots}{n_1! n_2! \dots n_k! \dots}. \quad (2.8)$$

The series of all Schur polynomials (of an increasing number of variables) is generated by

$$\exp \left( \sum_{k=1}^{\infty} t_k z^k \right) = \sum_{N=0}^{\infty} z^N S_N(t_1, t_2, \dots, t_k, \dots). \quad (2.9)$$

$J^l(z)$ , as computed from (2.5) (2.7), satisfy the recurrence relation

$$(l+1)J^l(z) = (J^0 + \partial)lJ^{l-1} = \dots = (J^0 + \partial)^l J^0 \quad (\equiv (l+1)!S_{l+1}(J^0, \dots), \quad \partial \equiv \frac{d}{dz}); \quad (2.10)$$

we have:  $2J^1 = (J^0)^2 + \partial J^0$ ,  $3J^2 = (J^0)^3 + 3J^0\partial J^0 + \partial^2 J^0$ ,  $4J^3 = (J^0)^4 + 6(J^0)^2\partial J^0 + 3(\partial J^0)^2 + 4J^0\partial^2 J^0 + \partial^3 J^0$  etc. (Viewed as quantum operators these expressions should be understood as normal products.) The local fields  $J^l$  (which, together with the unit operator span the infinite dimensional Lie algebra  $W_{1+\infty}$ ) are singled for their simple commutation relations ([BGT] Lemma 1.2) which are coded in the (singular terms of the ) OPE

$$\begin{aligned} J^k(z_1)J^l(z_2) &\simeq \sum_{\nu \geq 1} \left\{ \frac{l!}{(l-\nu)!} J^{k+l-\nu}(z_2) - (-1)^\nu \frac{k!}{(k-\nu)!} J^{k+l-\nu}(z_1) \right\} z_{12}^{-\nu-1} + \\ &+ (-1)^k l! k! z_{12}^{-k-l-2}. \end{aligned} \quad (2.11)$$

As we shall see shortly, the bilocal field (2.7) can also be expanded in a basis of local *quasiprimary* fields  $V^l(z)$  of dimension  $l+1$  which are expressed in

terms of  $J^l$  ([BGT] Lemma 1.3) by

$$V^l(z) = \frac{(l!)^2}{(2l)!} \sum_{k=0}^{\infty} (-1)^k \binom{l}{k} \binom{2l-k}{l} \partial^k J^{l-k}(z). \quad (2.12)$$

A characteristic feature of quasiprimary fields is their orthogonality; indeed, the 2-point function of a pair of  $V^l$ 's is

$$\langle V^l(z_1) V^{l'}(z_2) \rangle_0 = \frac{(l!)^2}{(2l)!} z_{12}^{-2l-2} \delta_{l,l'} \quad (2.13)$$

( $\langle \cdots \rangle_0$  being a shorthand for the vacuum expectation value  $\langle 0 | \cdots | 0 \rangle_0$ ).

According to [KR1] each positive energy unitary irreducible representation of  $W_{1+\infty}$  for  $c = 1$  is characterized by a single real number  $r$ , the  $J^0$  charge, and the existence of a unique ground state  $|r\rangle$  such that

$$(J_0^0 - r)|r\rangle = 0 = (L_0 - \frac{1}{2}r^2)|r\rangle, \quad J_n^0|r\rangle = 0 (= L_n|r\rangle) \text{ for } n > 0. \quad (2.14)$$

Here  $J_n^0$  are the  $J^0(z)$  Laurent modes, such that

$$J^0(z) = \sum_{n \in \mathbb{Z}} J_n^0 z^{-n-1}, \quad [J_n^0, J_{n'}^0] = n \delta_{n,-n'} \quad (2.15)$$

The *minimal quantum field theory* whose state space includes the irreducible  $W_{1+\infty}$  modules  $\mathcal{H}$  and  $\mathcal{H}_r$  (the *vacuum module* and the charge  $r \neq 0$  module  $\mathcal{H}_r$ ) is generated by a pair of oppositely charged fields  $\phi(z, \pm r)$ . Its OPE algebra contains (the  $W_{1+\infty}$  currents and) an infinite series of charged fields  $\phi(z, nr)$ ,  $n \in \mathbb{Z}$  (relatively local to  $J^0$  - see [BMT]). For  $r^2 = m \in \mathbb{N}$  they are also local among themselves Bose or Fermi fields; more precisely, they satisfy

$$z_{12}^{r_1 r_2} \phi(z_1, r_1) \phi(z_2, r_2) = z_{21}^{r_1 r_2} \phi(z_2, r_2) \phi(z_1, r_1) \\ \text{for } r_{1,2} \in \mathbb{Z} \sqrt{m} \text{ (} m = 1, 2, \dots \text{fixed)}. \quad (2.16)$$

In this case we recover the algebra  $\mathcal{A}_m$  generated by the fields  $\phi^\pm(z) = \phi(z, \pm \sqrt{m})$  satisfying (2.1), which provides an RCFT extension of  $W_{1+\infty}$ . Moreover, as noted in [BGT] (sec. 3) any RCFT extension of  $W_{1+\infty}$  for  $c = 1$  is of this type.

## 2.2 Expansion of the dipole field (2.1) into local quasiprimary fields $V^l(z, m)$

We introduce for arbitrary  $m$  ( $= 1, 2, \dots$ ) and  $c = 1$  the counterpart of (1.1), the *bilocal field*

$$V_m(z_1, z_2) = z_{12}^{m-1} \phi^+(z_1) \phi^-(z_2) - z_{12}^{-1} = \frac{1}{z_{12}} \left( : \exp \left\{ \int_{z_2}^{z_1} H(z) dz \right\} : - 1 \right); \quad (2.17)$$

it is *quasiprimary* of total dimension 1 in the sense that it satisfies

$$[L_n, V_m(z_1, z_2)] = \sum_{i=1}^2 z_i^n (z_i \frac{\partial}{\partial z_i} + \frac{n+1}{2}) V_m(z_1, z_2) \quad \text{for } n = 0, \pm 1. \quad (2.18)$$

(It is easily verified, in particular, that the differential operators in the right hand side annihilate, for  $n = 0, \pm 1$ , the 2-point function  $z_{12}^{-1}$ .) Eq. (2.17) (with the current  $H$  satisfying (2.2)) provides a version of the vertex operator construction, thus allowing to compute the correlation functions of  $V_m$ . We have, in particular,  $\langle V_m(z_1, z_2) \rangle_0 = 0$ ,

$$\begin{aligned} \langle V_m(z_1, z_2) V_m(z_3, z_4) \rangle_0 &= \frac{1}{z_{12} z_{34}} \left\{ (1 - \eta)^{-m} - 1 \right\} = \\ &= \frac{m}{z_{13} z_{24}} \sum_{n=0}^{\infty} \binom{m+n}{m} \frac{\eta^n}{n+1}, \quad \eta = \frac{z_{12} z_{34}}{z_{13} z_{24}}; \end{aligned} \quad (2.19)$$

$$\begin{aligned} \langle V_m(z_1, z_2) V_m(z_3, z_4) V_m(z_5, z_6) \rangle_0 &= (z_{12} z_{34} z_{56})^{-m} \left\{ 2 - \left( \frac{z_{13} z_{24}}{z_{23} z_{14}} \right)^m - \right. \\ &\quad \left. - \left( \frac{z_{35} z_{46}}{z_{36} z_{45}} \right)^m - \left( \frac{z_{15} z_{26}}{z_{16} z_{25}} \right)^m + \left( \frac{z_{13} z_{24} z_{35} z_{46} z_{15} z_{26}}{z_{23} z_{14} z_{36} z_{45} z_{16} z_{25}} \right)^m \right\}. \end{aligned} \quad (2.20)$$

We shall describe the complete OPE of (2.17) into local Bose fields thus extending to arbitrary  $m \in \mathbb{N}$  an earlier result about  $V_1$  (cf. Theorem 2.1 of [BGT]).

**Theorem 2.1** *The bilocal field (2.17) admits an expansion in integrals over local quasiprimary fields  $V^l$  (of dimension  $l+1$ ):*

$$V_m(z_1, z_2) = \sum_{l=0}^{\infty} \frac{z_{12}^l}{l!} \int_{z_2}^{z_1} p_l(z; z_1, z_2) V^l(z, m) dz \quad (2.21)$$

where  $p_l$  is the (normalized) weight function

$$p_l(z; z_1, z_2) = \frac{(2l+1)!}{(l!)^2 z_{12}^{2l+1}} (z_1 - z)^l (z - z_2)^l \left( \int_{z_2}^{z_1} p_l(z; z_1, z_2) dz = 1 \right). \quad (2.22)$$

The 2-point functions of  $V^l$  (which display their orthogonality),

$$\frac{l+1}{l!} \langle V^l(z_1, m) V^{l'}(z_2, m) \rangle_0 = C_{lm} z_{12}^{-2l-2} \delta_{l,l'}, \quad (2.23)$$

involve the same normalization constant as the 3-point function

$$(l+1)\langle V_m(z_1, z_2)V^l(z_3, m)\rangle_0 = C_{lm}\frac{z_{12}^l}{(z_{13}z_{23})^{l+1}}. \quad (2.24)$$

The (positive) constants  $C_{lm}$  are uniquely determined by the 4-point function (2.19). Conversely,  $V^l(z, m)$  are computed from  $V_m(z_1, z_2)$  by differentiation:

$$V^l(z, m) = \frac{l!}{(2l)!} \lim_{z_1, z_2 \rightarrow z} \left\{ \partial_1^l (-\partial_2)^l \left( z_{12}^l V_m(z_1, z_2) \right) \right\}. \quad (2.25)$$

Furthermore,  $V^l$  are again related to the Schur polynomials  $S_n(H, \frac{\partial H}{2!}, \dots, \frac{\partial^{n-1} H}{n!})$  by (the counterpart of) Eq. (2.12).

The **proof** follows, for the most part the argument in [BGT] and we shall only sketch it emphasizing the new points.

The weight function (2.22) satisfies the identity

$$\int_{z_2}^{z_1} p_l(z; z_1, z_2)(z - z_3)^{-2l-2} dz = z_{13}^{-l-1} z_{23}^{-l-1}. \quad (2.26)$$

Since, on the other hand, 2- and 3- point functions of quasiprimary fields are determined (up to normalization constants) by Möbius (conformal) invariance, this implies

$$\langle V_m(z_1, z_2)V^l(z_3, m)\rangle_0 = \frac{z_{12}^l}{l!} \int_{z_2}^{z_1} p_l(z; z_1, z_2) \langle V^l(z, m)V^l(z_3, m)\rangle_0 dz. \quad (2.27)$$

Orthogonality of  $V^l$  (reflected in (2.23)) is a general property of quasiprimary fields of different dimensions. Inserting the expansion (2.21) for  $V_m(z_3, z_4)$  into (2.19) and using (2.24) we end up with a triangular set of linear equations for  $C_{lm}$ :

$$(1 - \eta)^{-m} = 1 + \sum_{l=0}^{\infty} C_{lm} \frac{\eta^{l+1}}{(l+1)!} F(l+1, l+1; 2l+2; \eta). \quad (2.28)$$

Here we have used the following representation for the Gauss hypergeometric function

$$F(n, n; 2n; \eta) = z_{13}^n z_{24}^n \int_{z_2}^{z_1} \frac{p_l(z; z_1, z_2) dz}{(z_1 - z)^n (z_2 - z)^n} \left( \eta = \frac{z_{12} z_{34}}{z_{13} z_{24}} \right). \quad (2.29)$$

We thus have

$$\sum_{l=0}^{\infty} C_{lm} \frac{\eta^{l+1}}{(l+1)!} \sum_{n=0}^{\infty} \binom{l+n}{n}^2 \binom{2l+1+n}{n}^{-1} \eta^n = \sum_{l=0}^{\infty} \binom{m+l}{l+1} \eta^{l+1} \quad (2.30)$$

The first seven  $C_{lm}$  are captured by the formula

$$C_{lm} = m^{l+1} + \binom{l+1}{4} \frac{m^{l-1}}{2l-1} + \binom{l+1}{6} \frac{5l^2 - 7l + 6}{8(2l-1)(2l-3)} m^{l-3} \quad \text{for } 0 \leq l \leq 6. \quad (2.31)$$

On the other hand, for  $m = 1$ , we recover the expression  $C_{l1} = (l+1)! \binom{2l}{l}^{-1}$  of [BGT]. The important point is the existence of a unique solution of the (infinite) system (2.30)  $C_{lm} > 0$ .

Finally, Lemma 1.3 of [BGT] holds in this case, too, and we can write

$$V^l(z, m) = \frac{(l!)^2}{(2l)!} \sum_{k=0}^l (-1)^k \binom{l}{k} \binom{2l-k}{l} (l-k)! \partial^k S_{l-k+1}(H, \frac{\partial H}{2!}, \dots, \frac{\partial^{l-k} H}{(l-k+1)!}) \quad (2.32)$$

where  $S_{n+1}(H, \frac{\partial H}{2!}, \dots, \frac{\partial^n H}{(n+1)!}) = (H + \partial)^n H$  (cf. Eq. (2.10)) and we omit (here and in the following remark) the normal product sign.

**Remark 2.1.** In spite of the fact that  $V^l(z) = V^l(z, 1)$  is expressed by the same Schur polynomials as  $V^l(z, m)$  - only the argument  $H$  is substituted by the normalized current  $J^0$  in  $V^l(z, 1)$  - the quasiprimary fields  $V^l(z, m)$  do not close under commutation among themselves (and the unit operator) for  $m \geq 2$ . In effect the fields  $V^l(z, m) = m^{\frac{l+1}{2}} V^l(z)$  for  $l = 0, 1, 2$  generate the whole  $W_{1+\infty}$  algebra spanned by  $V^l(z)$  (and 1). On the other hand, the fields  $V^l(z, m)$  and  $V^l(z)$  are linearly independent for  $l \geq 3$ ,  $m \geq 2$ . Indeed, we have

$$\begin{aligned} V^3(z, m) &= \frac{H^4(z)}{4} + \frac{1}{20} P_1(H, H', H'') = \frac{m^2}{4} J^4(z) + \frac{m}{20} (2JJ'' - 3J'^2)(z) \\ (\text{for } H = \sqrt{m}J); \quad V^4(z, m) &= \frac{1}{5} H^5 + \frac{1}{7} H P_1, \text{ etc.} \end{aligned} \quad (2.33)$$

Here  $P_1$  is the first of a series of quasiprimary fields that are quadratic forms  $P_n$  of  $H$  and its derivatives (each term having dimension  $2n + 2$ ):

$$\begin{aligned} P_n(H, H', \dots, H^{(2n)}) &= 2 \sum_{l=0}^{n-1} (-1)^l a_l H^{(l)} H^{(2n-l)} + (-1)^n a_n (H^{(n)})^2, \\ a_l &= \prod_{k=0}^{l-1} \binom{2n-1+k}{2} \prod_{k=l}^{n-1} \binom{k+2}{2}. \end{aligned} \quad (2.34)$$



### 3 Extension to arbitrary $c \in \mathbb{N}$ . Discussion

#### 3.1 Additive bilocal fields and lattice RCFT extensions

The construction of Sec. 2 extends to arbitrary positive integer  $c$  due to the following simple observation.

**Proposition 3.1** ([BGT], Lemma 2.2) *Let  $V_i(z_1, z_2)$ ,  $1 \leq i \leq c$  be a commuting set of bilocal fields with the properties of  $V_1(z_1, z_2)$  (2.7). Then their sum generates the  $W_{1+\infty}$  algebra for central charge  $c$ :*

$$\begin{aligned} \sum_{i=1}^c V_i(z_1, z_2) &= \sum_{l=0}^{\infty} \frac{z_{12}^l}{l!} J^l(z_2) \Rightarrow \\ [J^k(z_1), J^l(z_2)] &= \sum_{\nu \geq 1} \left\{ (-1)^\nu \binom{l}{\nu} J^{k+l-\nu}(z_2) - \binom{k}{\nu} J^{k+l-\nu}(z_1) \right\} \delta^{(\nu)}(z_{12}) \Big\} - \\ &- (-1)^l \frac{k!l!c}{(k+l+1)!} \delta^{(k+l+1)}(z_{12}). \end{aligned} \quad (3.1)$$

If we use the bosonized expressions

$$V_i(z_1, z_2) = \frac{1}{z_{12}} \left( : \exp \left\{ \int_{z_2}^{z_1} J_i^0(z) dz \right\} : - 1 \right) \quad \text{with} \quad [J_i^0(z_1), J_j^0(z_2)] = -\delta'(z_{12}) \delta_{ij}, \quad (3.2)$$

then  $J^l(z) = \sum_{i=1}^c J_i^l(z)$  where  $J_i^l$  is expressed as a Schur polynomial of  $J_i^0(z)$  and its derivatives

$$J_i^l(z) = l! S_{l+1}(J_i^0(z), \frac{\partial J_i^0(z)}{2!}, \dots, \frac{\partial^l J_i^0(z)}{(l+1)!}). \quad (3.3)$$

Similarly, the basic quasiprimary fields  $V^l(z)$  appear as sums of mutually commuting  $c = 1$  fields  $V_i^l(z, 1)$ .

Let  $W^{(c)}$  be the chiral algebra corresponding to the vacuum representation of  $W_{1+\infty}$  of central charge  $c$ ; it is spanned by (normal ordered) polynomials of  $V^l(z)$ . The set of RCFT extensions is much richer for  $c > 1$ . It includes, in particular, local charged fields' algebras  $\mathcal{A}(L)$  associated with a  $c$ -dimensional integral Euclidean charge lattice  $L$ . In other words, there is a basis  $(\mathbf{q}^1, \dots, \mathbf{q}^c)$  in the space of  $U(1)^{\times c}$  charges such that the (symmetric) positive definite Gram matrix  $(\mathbf{q}^i | \mathbf{q}^j)$  of inner products has integer entries. To each such lattice corresponds a vertex algebra (for a recent review - see

[K]) spanned by charged vertex operators  $Y(\mathbf{q}, z)$  and “Cartan currents”  $H^{\mathbf{q}}(z)$  satisfying OPE of the type

$$Y(\mathbf{q}, z_1)Y(-\mathbf{q}, z_2) = \frac{1}{z_{12}^{|\mathbf{q}|^2}} \exp \left\{ \int_{z_2}^{z_1} H^{\mathbf{q}}(z) dz \right\}, \quad H^{\mathbf{q}}(z_1)H^{\mathbf{q}'}(z_2) \sim \frac{(\mathbf{q}|\mathbf{q}')}{z_{12}^2}, \quad (3.4)$$

$$Y(\mathbf{q}, z_1)Y(\mathbf{q}', z_2) \sim \varepsilon(\mathbf{q}, \mathbf{q}') z_{12}^{(\mathbf{q}|\mathbf{q}')} Y(\mathbf{q} + \mathbf{q}', z_2) \quad (\text{for } \mathbf{q} + \mathbf{q}' \neq 0) \quad (3.5)$$

where  $\varepsilon$  is a  $(\pm 1)$  valued cocycle that ensures the correct spin-statistics relation between different vertex operators. If the charge lattice is odd, -i.e., if there are vectors  $\mathbf{q} \in L$  such that  $|\mathbf{q}|^2 \equiv (\mathbf{q}|\mathbf{q})$  is odd (which is the case of interest in applications to a quantum Hall system - see [FST] [CTZ] [CZ] and references therein) then the resulting chiral algebra  $\mathcal{A}(L)$  is a local  $\mathbb{Z}_2$  graded superalgebra. In order to identify a  $W^{(c)}$  subalgebra we should choose a 1-dimensional subspace in the  $c$ -dimensional space of currents, thus singling out the current  $J^0(z) = \sum_{i=1}^c J_i^0(z)$ . This is achieved, in particular, by selecting an “electric charge” vector  $\mathbf{Q}$  in the dual lattice  $L^*$  (whose square  $|\mathbf{Q}|^2$  gives the quantum Hall filling fraction [FST] [CTZ]).

The positive energy representations of the RCFT so obtained, in which all vertex operators  $Y(\mathbf{q}, z)$ ,  $\mathbf{q} \in L$  are single valued, are in one-to-one correspondence with the elements of the (finite) abelian group  $L^*/L$  of order

$$|L^*/L| = \det \left( (\mathbf{q}^i | \mathbf{q}^j) \right).$$

Let  $\Gamma$  be any finite group of automorphisms of the chiral algebra  $\mathcal{A}(L)$  which leaves each element of its  $W^{(c)}$  subalgebra invariant. (It is enough to demand the invariance of the first three “tensor currents”  $V^l(z)$ ,  $l = 0, 1, 2$  which generate the entire  $W_{1+\infty}$  algebra.) There is a wealth of such inner automorphism groups in the case when  $L$  is a semisimple Lie group lattice (for integer  $c$  it may only corresponds to a lattice of the A-D-E type, the basis  $\mathbf{q}^i$  being identified with a basis of simple roots). Viewing  $\Gamma$  as a *gauge group* we define the *observable algebra* as the subalgebra  $\mathcal{A}(L)^\Gamma$  of  $\Gamma$  invariant elements of  $\mathcal{A}(L)$ . The resulting chiral algebra  $\mathcal{A}(L)^\Gamma$  provides another RCFT extensions of  $W^{(c)}$  (see [KT]).

The characters of positive energy irreducible representations of a lattice current algebra span, in general, a finite dimensional representation of the subgroup of the modular group  $\text{SL}(2, \mathbb{Z})$  generated by the matrices

$$S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad \text{and} \quad T^2 = \begin{pmatrix} 1 & 2 \\ 0 & 1 \end{pmatrix} \quad (3.6)$$

(If  $L$  is an even lattice then they span a representation of the full  $SL(2, \mathbb{Z})$ .) Recently, all weak modular invariants (i.e. invariants under the group generated by (3.6) have been classified -see [G] ). On the other hand, given a *non-exceptional gauge group* of inner automorphisms  $\Gamma$  (such is, in particular, any finite subgroup of  $U(n)$  or  $SU(n)$  ), it has been demonstrated in [KT] how to construct the specialized characters (that only capture the energy distribution) of the corresponding  $\mathcal{A}(L)^\Gamma$  *orbifold modules* and to exhibit their modular transformation properties.

**Remark 3.1.** Albeit we thus arrive at a rich family of manageable RCFT extensions of  $W_{1+\infty}$ , the set of such extensions is not exhaustive - even if we admit outer automorphisms in  $\Gamma$ , for which no general construction of orbifold modules is known. RCFT extensions of a different type are obtained through the so called conformal embedding ([SW],[BN],[AGO],[S],[T],[RST]). A simple example of this type is provided by the level 5  $su(3)$ -current algebra  $\mathcal{A}_5(su(3))$  of Virasoro central charge  $c_k = \frac{8k}{3+k} = 5$  for  $k = 5$ , which can be viewed as the *observable subalgebra* of the rank 5 lattice chiral algebra  $\mathcal{A}_1(su(6))$ . Indeed, the vacuum space of  $\mathcal{A}_1(su(6))$  splits into a direct sum of two  $\mathcal{A}_5(su(3))$  modules (with the same central charge  $c = 5$ , and hence, the same Virasoro subalgebra): one with  $su(3)$  weight  $\Lambda_0 = (0, 0)$  and ground state energy  $\Delta_0 = 0$  and a second one with  $\Lambda_1 = (2, 2)$  and a degenerate 27 dimensional ground state of energy (=conformal weight)  $\Delta_1 = \frac{1}{24} \left\{ \lambda_1(\lambda_1 + 3) + \lambda_2(\lambda_2 + 3) + \lambda_1 \lambda_2 \right\} = 1$  for  $\lambda_1 = \lambda_2 = 2$ . Thus  $\mathcal{A}_5(su(3))$  can be viewed as an RCFT extension of either  $W^{(5)}$  or the part of  $W^{(6)}$  orthogonal to the  $u(1)$  current  $J^0$ . It can not be viewed as the gauge invariant part of the lattice algebra  $\mathcal{A}_1(su(6))$  with respect to a finite gauge group  $\Gamma$  since then the  $\mathcal{A}_5(su(3))$  module of highest weight  $\Lambda_1$  should have an integer quantum dimension (equal to the dimension of the associated irreducible representation of  $\Gamma$ -see [RST] and references therein) while it is, in fact, irrational:

$$d_q(\Lambda) = \frac{[3]^2[6]}{[2]} = (1 + \sqrt{2})^2 \quad \text{for } q^8 = -1.$$

Only for  $c = 1$  we have listed all RCFT extensions of the  $u(1)$  current algebra - cf.[PT] -and hence all extensions of  $W^{(1)}$ .

### 3.2 Computing character dependence on chemical potential for orbifold models

It is important, as far applications to the quantum Hall effect are concerned, to compute the character dependence on the chemical and electric potentials

combined in a complex variable  $\zeta$  (see [CZ]; the  $\zeta$  dependence carries information about charge distribution and the so called spectral flow). In the case of lattice orbifolds this requires the following simple (but important) complement of the algorithm presented in [KT].

To set the stage we reformulate briefly some notions and results of [KT] adapted to the present context.

Let  $L$  be an integral euclidean lattice and let  $\mathbf{Q}$  be a fixed vector in the dual lattice  $L^*$ , the “electric charge” (such a pair  $(L, \mathbf{Q})$ , satisfying some additional requirements, is called a *chiral quantum Hall lattice* in [FST]). We shall assume, for the sake of convenience, that  $L$  is *maximal*, -i.e., there is no sublattice of  $L^*$  that is a proper extension of  $L$ . (This assumption does not restrict the generality of the orbifold chiral algebra we shall end up with.) Let  $G$  be a connected compact Lie group whose Lie algebra is generated by the 0-components of all (dimension 1) currents in the chiral algebra  $\mathcal{A}(L)$ . It can be hence assumed acting by unitary operators in each irreducible (positive energy)  $\mathcal{A}(L)$  module.  $G$  is a rank  $l = \dim L$  Lie group: it has a maximal abelian subgroup  $U(1)^{\times l}$  where  $l = \text{rank } L$ ; the center  $\mathbb{Z}_G \subset U(1)^{\times l}$  of  $G$  is nontrivial: it contains a (finite) subgroup isomorphic to  $|L^*/L|$ . To any  $\mathbf{q} \in L^*$  there corresponds a pair of charged vertex operators  $Y(\pm \mathbf{q}, z)$  (which intertwine the vacuum representation of  $\mathcal{A}(L)$  with the charge  $\pm \mathbf{q}$  representations) and hence a  $u(1)$  current  $H^{\mathbf{q}}(z)$  satisfying (3.4) whose 0-component  $H_0^{\mathbf{q}}$  belongs to the Lie algebra  $\mathcal{G}$  of  $G$ . ( $\mathcal{G}$  may also involve raising and lowering operators  $E^{\pm \underline{\alpha}}$  for  $\underline{\alpha} \in L$ ,  $|\underline{\alpha}|^2 = 2$  whenever  $L$  contains such vectors  $\underline{\alpha}$ . Let  $\Gamma$  be a (non-exceptional) finite subgroup of  $G$  acting by inner automorphisms on  $\mathcal{A}(L)$  and leaving the current  $H^{\mathbf{Q}}(z)$  invariant. To each inner automorphism  $\rho \in \text{Ad } \Gamma$  of  $\mathcal{A}(L)$  we make correspond a  $\underline{\beta} = \underline{\beta}_\rho \in i\mathcal{G}$  such that

$$\rho(V) = b V b^{-1}, \quad b = e^{2\pi i \underline{\beta}}, \quad \Gamma_b = \Gamma_{\underline{\beta}} \quad (3.7)$$

where  $\Gamma_b(\Gamma_{\underline{\beta}})$  is the centralizer of  $b(\underline{\beta})$  in  $\Gamma$ . (We identify vectors  $\underline{\beta} \in L$  with corresponding Lie algebra element  $H_0^{\underline{\beta}}$ .)

Let  $\mathcal{H}_{\underline{\lambda}}$ ,  $\underline{\lambda} \in L^*/L$  be an irreducible positive energy  $\mathcal{A}(L)$  module. Define its character  $\chi_{\underline{\lambda}}$

$$\chi_{\underline{\lambda}}(\tau, \underline{\alpha}) = \text{tr}_{\mathcal{H}_{\underline{\lambda}}}(q^{L_0 - \frac{c}{24}} e^{2\pi i \underline{\alpha}}), \quad q = e^{2\pi i \tau}, \quad \text{Im } \tau > 0; \quad i\underline{\alpha} \in \mathcal{G} \quad (3.8)$$

It follows that

$$\chi_{\underline{\lambda}}(\tau, \underline{\alpha}) = \frac{1}{[\eta(\tau)]^l} \sum_{\underline{\gamma} \in L + \underline{\lambda}} q^{\frac{1}{2}(\underline{\gamma}|\underline{\gamma})} e^{2\pi i (\underline{\gamma}|\underline{\alpha})} \quad (\eta = q^{\frac{1}{24}} \prod_{n=1}^{\infty} (1 - q^n), \quad l = \text{rank } L). \quad (3.9)$$

The character of a  $\underline{\beta}$ -twisted module  $\mathcal{H}_{\underline{\lambda}}^{\underline{\beta}}$  is (see Eq. (3.18) of [KT])

$$\chi_{\underline{\lambda}}^{\underline{\alpha}, \underline{\beta}}(\tau) = q^{\frac{1}{2}|\underline{\beta}|^2} \chi_{\underline{\lambda}}(\tau, \underline{\alpha} - \tau \underline{\beta}), \quad [\underline{\alpha}, \underline{\beta}] = 0. \quad (3.10)$$

**Proposition 3.2** *Let  $\sigma$  be (the character of) an irreducible representation of the centralizer  $\Gamma_{\underline{\beta}} (= \Gamma_b)$  of  $\underline{\beta}$  in  $\Gamma$ . Then the projected module  $\mathcal{H}_{\underline{\lambda}\sigma}^{\underline{\beta}}$  is only nontrivial if  $\sigma$  and  $\underline{\lambda}$  agree on  $L^*/L$  and then the character is*

$$\chi_{\underline{\lambda}\sigma}^{\underline{\beta}}(\tau, \zeta) = \frac{1}{|\Gamma_{\underline{\beta}}|} \sum_{h=e^{2\pi i \underline{\alpha}} \in \Gamma_{\underline{\beta}}} e^{-2\pi i \zeta(\mathbf{Q}|\underline{\beta})} \chi_{\underline{\lambda}}^{\zeta \mathbf{Q} + \underline{\alpha}, \underline{\beta}}(\tau) \sigma^*(h) \quad (3.11)$$

$|\Gamma_{\underline{\beta}}|$  standing for the number of elements of  $\Gamma_{\underline{\beta}}$ .

Indeed, defining  $\mathcal{H}_{\underline{\lambda}\sigma}^{\underline{\beta}}$  by

$$\mathcal{H}_{\underline{\lambda}\sigma}^{\underline{\beta}} = P_{\sigma} \mathcal{H}_{\underline{\lambda}}^{\underline{\beta}} \text{ where } P_{\sigma} = \frac{\sigma(1)}{|\Gamma_{\underline{\beta}}|} \sum_{h \in \Gamma_{\underline{\beta}}} \sigma^*(h) h (= P_{\sigma}^2) \quad (3.12)$$

one proves by a simple extension of the argument of Sec. 4.1 of [KT] that

$$\chi_{\underline{\lambda}\sigma}^{\underline{\beta}}(\tau, \zeta) = \text{tr}_{\mathcal{H}_{\underline{\lambda}\sigma}^{\underline{\beta}}} (q^{L_0 - H_0^{\underline{\beta}} + \frac{1}{2}|\underline{\beta}|^2 - \frac{c}{24}} e^{2\pi i \zeta(H_0^{\underline{\beta}} - (\mathbf{Q}|\underline{\beta}))}). \quad (3.13)$$

Instead of reproducing the details we shall work out an example of physical interest (cf. [CGT]).

**Example 3.1.** Let  $L$  be a 2-dimensional orthogonal lattice with basis  $\mathbf{e}^1, \mathbf{e}^2$  and Gram matrix

$$\left( (\mathbf{e}^i | \mathbf{e}^j) \right) = \begin{pmatrix} (\mathbf{e}^1 | \mathbf{e}^1) & (\mathbf{e}^1 | \mathbf{e}^2) \\ (\mathbf{e}^2 | \mathbf{e}^1) & (\mathbf{e}^2 | \mathbf{e}^2) \end{pmatrix} = \begin{pmatrix} m & 0 \\ 0 & 1 \end{pmatrix}, \quad m \in \mathbb{N}, \quad (3.14)$$

and let  $\mathbf{Q} = \mathbf{e}_1^*$  (where  $\{\mathbf{e}_i^*\}$  stands for the dual basis in  $L^*$  satisfying  $(\mathbf{e}_i^* | \mathbf{e}^j) = \delta_j^i$ ). Consider a  $\mathbb{Z}_M$  gauge group in  $\mathcal{A}(L)$  generated by an automorphism of type (3.7) with

$$\underline{\beta} = \underline{\beta}_M = \frac{1}{M} (\mu \mathbf{e}_1^* + \mathbf{e}_2^*) \quad 0 < \mu < M. \quad (3.15)$$

$\mathcal{A}(L)$  has  $m$  (untwisted) representations with partition functions

$$\chi_l(\tau, \underline{\alpha}) = K_l(\tau, \alpha^1; m) K_0(\tau, \alpha^2; 1) \quad \text{for } \underline{\alpha} = \alpha^i \mathbf{e}_i^*, \quad (3.16)$$

where  $K_l$  are the 1-dimensional lattice characters (see, e.g., [PT])

$$K_l(\tau, \alpha; m) = \frac{1}{\eta(\tau)} \sum_{n \in \mathbb{Z}} q^{\frac{m}{2}(n + \frac{l}{m})^2} e^{2\pi i \alpha(n + \frac{l}{m})} \quad (3.17)$$

$l \bmod m$  (for  $m$  even a canonical choice is  $1 - \frac{m}{2} \leq l \leq \frac{m}{2}$ ).

The lift  $b = e^{2\pi i H_0^\beta}$  ( $H_0^\beta = \frac{1}{M}(\mu H_0^{\mathbf{e}_1^*} + H_0^{\mathbf{e}_2^*})$ ) of the automorphism group  $\mathbb{Z}_M$  to the  $\mathcal{A}(L)$  modules  $\mathcal{H}_l$  is  $\mathbb{Z}_{mM}$ . Setting

$$\begin{aligned} \sigma^*(e^{2\pi i \underline{\alpha}}) &= e^{-2\pi i \frac{\underline{\alpha}}{m} \sigma} \text{ (for } \underline{\alpha} = \alpha(\mu \mathbf{e}_1^* + \mathbf{e}_2^*)\text{),} \\ \underline{\beta} &= \beta(\mu \mathbf{e}_1^* + \mathbf{e}_2^*); \quad e^{2\pi i \frac{\underline{\alpha}}{m}}, e^{2\pi i \frac{\underline{\beta}}{m}} \in \mathbb{Z}_{mM}, \quad |\Gamma_{\underline{\beta}}| = mM \end{aligned} \quad (3.18)$$

we can evaluate the finite sum in (3.11) with the result

$$\chi_{l,\sigma}^\beta(\tau, \zeta) = \frac{1}{\eta^2(\tau)} \sum_{n_1, n_2 \in \mathbb{Z}} e^{2\pi i(n_1 + \frac{l - \mu \beta}{m})\zeta} q^{\frac{m}{2}(n_1 + \frac{l - \mu \beta}{m})^2 + \frac{1}{2}(n_2 - \beta)} \delta_{m(\mu n_1 + n_2) + \mu l}^{\sigma \bmod mM}. \quad (3.19)$$

It follows that  $\sigma - \mu l = 0 \bmod m$ , in accord with the first statement in Proposition 3.1, and we can take the Kronecker  $\delta$  into account setting (for  $M, \mu$  coprime)

$$\begin{aligned} n_1 &= \frac{\sigma - \mu l}{m} + r + Mn, \quad n_2 = \frac{\mu l - \sigma}{m}(\mu - 1) - \mu r + mn', \\ r &= 0, 1, \dots, M-1 \quad (n, n' \in \mathbb{Z}). \end{aligned} \quad (3.20)$$

Inserting in (3.19) and using once more (3.17) we find

$$\begin{aligned} \chi_{l,\sigma}^{\nu/M}(\tau, \zeta) &= \sum_{r=0}^{M-1} K_{M[\sigma - (\mu-1)l] - \mu\nu + mM r}(\tau, M\zeta; mM^2) \times \\ &\quad \times K_{M \frac{l\mu - \sigma}{m}(\mu-1) - \nu - \mu M r}(\tau, 0; M^2) \end{aligned} \quad (3.21)$$

**Remark 3.2.** A non-analytic factor,  $e^{-\frac{\pi}{m} \frac{(Im \zeta)^2}{Im \tau}}$ , is to be assigned to the character in order to ensure its  $V : \zeta \rightarrow \zeta + \tau$  covariance (see [CZ]).

**Remark 3.3.** Given the automorphism  $\rho$  there is a “gauge” freedom in the choice of  $b$  and  $\underline{\beta}$  in (3.7) (in the example (3.15) it amounts to adding to  $\underline{\beta}$  an arbitrary vector of  $L^*$ ). Theorem 4.2 of [KT] guarantees that the resulting orbifold models do not depend on this choice.

*To sum up:* quantum field theory realization of  $W_{1+\infty}$  are provided by  $U(1)^{\times c}$  current algebras which also allow to construct a large family of RCFT

extensions of such algebras. These leave room for additional physical requirements having in mind specific applications, - e.g., to quantum Hall systems.

The present paper is an outgrow of previous work in collaboration with Bojko Bakalov to whom we are indebted, in particular, for an indication of the relevance of Schur's polynomials in the present context. The work has been supported in part by the Bulgarian Foundation for Scientific Research under contract F-404.

## References

- [AFMO] H. Awata, M. Fukuma, Y. Matsuo, S. Odake, *Representation theory of the  $W_{1+\infty}$  algebra*, Prog. Theor. Phys.; Proc. Suppl. **118** (1995) 343-373, (hep-th/9408158).
- [AGO] J.R.C. Arcari, J.F. Gomez, D.I. Olive, Nucl. Phys. **B285** (1987) 327.
- [BGT] B.N. Bakalov, L.S. Georgiev, I.T. Todorov, *A QFT approach to  $W_{1+\infty}$* , in **New Trends in Quantum Field Theory** Proc. of 1995 Razlog (Bulgaria) Workshop, A. Ganchev et al. (eds.) (Heron Press, Sofia 1996), pp. 147 – 158 (hep-th/9512078).
- [BHY] B. Bakalov, E. Horozov, M. Yakimov, *Highest weight modules over  $W_{1+\infty}$  algebra and the bispectral problem*, Sofia preprint, (q-alg/9602012).
- [BMT] D. Buchholz, G. Mack, I.T. Todorov, Nucl. Phys. B (Proc. Suppl.) **5B** (1988) 20.
- [BN] P. Bouwknegt, W. Nahm, Phys. Lett. **B184** (1987) 359.
- [CGT] A. Cappelli, L.S. Georgiev, I.T. Todorov, (in preparation).
- [CTZ] A. Cappelli, C. A. Trugenberger, G. R. Zemba, Nucl. Phys. **B448** (1995) 470.
- [CZ] A. Cappelli, G.R. Zemba, *Modular invariant partition functions in the quantum Hall effect*, hep-th/9605127.
- [DFZ] G.F. Dell' Antonio, V. Frishman, D. Zwanziger, Thirring model in terms of currents: solution and light-cone expansion, Phys. Rev. **D6** (1972) 988-1007.
- [FKRW] E. Frenkel, V. Kac, A. Radul, W. Wang,  $W_{1+\infty}$  and  $W(gl_N)$  with central charge  $N$ , Commun. Math. Phys. **170** (1995) 337-357.

- [FST] J. Fröhlich, M. Thiran, J. Stat. Phys. **76** (1994) 209; J. Fröhlich, U.M. Studer, M. Thiran, *A classification of quantum Hall fluids*, cond-mat/9503113.
- [G] T. Gannon,  *$U(1)^m$  modular invariants,  $N = 2$  minimal models, and the quantum Hall effect*, hep-th/9608063, Nucl. Phys. B
- [K] V. G. Kac, *Vertex algebras*, in **New Trends in Quantum Field Theory** Proc. of 1995 Razlog (Bulgaria) Workshop, A. Ganchev et al. (eds.) (Heron Press, Sofia 1996), pp. 261-358; **Vertex Algebras for Beginners** (AMS University Lecture Series, Vol. 10, 1997) 141 p.
- [KR1] V. Kac, A. Radul, Quasifinite highest weight modules over the Lie algebra of differential operators on the circle, Commun. Math. Phys. **157** (1993) 429–457 (hep-th/9308153).
- [KR2] V. Kac, A. Radul, Representation theory of the vertex algebra  $W_{1+\infty}$ , Transformation Groups **1** (1996) 41-70 (hep-th/9512150).
- [KR3] V. Kac, A.K. Raina, Bombay Lectures on Highest Weight Representations of Infinite Dimensional Lie Algebras, Advanced Ser. in Math. Phys. **2** (World Scientific,
- [KT] V.G. Kac and I.T. Todorov, *Affine orbifolds and rational conformal field theory extensions of  $W_{1+\infty}$* , hep-th/9612078v2, 15 Dec. 1996, Commun. Math. Phys. (to appear).
- [PT] R. R. Paunov, I. T. Todorov, Phys. Lett. **B196** (1987) 519–526.
- [RST] K.-H. Rehren, Ya.S. Stanev, I.T. Todorov, Commun. Math. Phys. **174** (1996) 605.
- [S] Ya.S. Stanev, J. Math. Phys. **36** (1995) 2057.
- [SW] A.N. Schellekens, N.P. Warner, Phys. Rev. **D34** (1986) 3092.
- [T] I. T. Todorov, What are we learning from 2-dimensional conformal models?, in: Mathematical Physics Towards the 21st Century, R.N. Sen and A. Gersten eds. (ben-Gurion Univ. of the Negev Press, Beer-Sheva 1994) pp. 160-176.